



Rabeprazole & Domperidone

Rabappi-DSR

PRODUCT INFORMATION LEAFLET

Brand Name: **[Rabappi-DSR]**

Generic Name: Rabeprazole and Domperidone capsules

Strength: Rabeprazole 20 mg (EC) + Domperidone 30 mg (SR)

Dosage Form: capsules

COMPOSITION

Each hard gelatin capsule contains:

- Rabeprazole Sodium IP equivalent to Rabeprazole ... 20 mg (Enteric Coated)
- Domperidone Maleate IP equivalent to Domperidone ... 30 mg (Sustained

Release)

Excipients: q.s.

DESCRIPTION

Rabeprazole is a proton pump inhibitor (PPI) that suppresses gastric acid secretion by inhibiting the H⁺/K⁺ ATPase enzyme in the stomach lining.

Domperidone is a prokinetic and antiemetic that enhances gastrointestinal motility and prevents nausea and vomiting.

This combination is effective in the treatment of acid-related gastrointestinal disorders and improves gastric emptying.

INDICATIONS

Used in the treatment of:

- Gastroesophageal reflux disease (GERD)
- Acid reflux and heartburn
- Functional dyspepsia
- Non-ulcer dyspepsia



- Nausea and vomiting related to acid peptic disorders
- Gastric and duodenal ulcers

DOSAGE AND ADMINISTRATION

Adults: One capsule daily, preferably 15–30 minutes before a meal, or as directed by the physician.

Do not chew or crush.

For best results, take at the same time each day.

CONTRAINDICATIONS

- Known hypersensitivity to rabeprazole, domperidone, or any component of the formulation
- Patients with prolactin-secreting tumors (prolactinoma)
- Patients with gastrointestinal hemorrhage, mechanical obstruction, or perforation
- Domperidone is contraindicated in patients with moderate to severe hepatic impairment or prolonged QT interval

PRECAUTIONS

- Use with caution in patients with liver or kidney dysfunction
- Long-term use of rabeprazole may result in vitamin B12 deficiency, hypomagnesemia, or osteoporosis-related fractures
- Domperidone may cause QT prolongation; monitor ECG and electrolytes in at-risk patients
- Avoid concurrent use with other QT-prolonging medications

SIDE EFFECTS

Common:

- Headache
- Diarrhea or constipation
- Abdominal pain
- Dizziness
- Dry mouth

Rare:



- Hypersensitivity reactions (rash, itching)
- Elevated liver enzymes
- Galactorrhea, gynecomastia (due to domperidone)
- QT interval prolongation

STORAGE INSTRUCTIONS

Store in a cool, dry place below 25°C.
Protect from light and moisture.
Keep out of reach of children.

PRESENTATION

Box of 10x10 capsules in Aluminium strip pack.

SCHEDULE

Schedule H Prescription Drug – Not to be sold without the prescription of a registered medical practitioner.

Made in India for:



H.no:1-2, GaneshNagarColony, Kuntloor, Hayathnagar
R.R.Dist, Telangana, 501505.